


TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		
International Application. No.	International Filing Date	U.S. Application No. 09/869106 Unassigned
PCT/FR99/03270	December 23, 1999	Priority Date Claimed December 24, 1998




Title of Invention  
**GRAM-POSITIVE BACTERIA DEPRIVED OF HtrA PROTEASE ACTIVITY AND THEIR USES**  
 Applicants For DO/EO/US  
**Isabelle POQUET, Alexandra GRUSS, Alexandre BOLOTINE and Alexei SOROKINE**

Applicants herewith submit to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(4)).
9. ☐ An oath or declaration of the inventors (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 14. below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
14. ☒ A SECOND or SUBSEQUENT preliminary amendment.
14. ☒ Other items or information:
  - a. ☒ WO 00/39309
  - b. ☒ PCT/IB/304
  - ☒ PCT/IB/308
  - ☒ PCT/IPEA/409
  - ☒ Paper Copy of Sequence Listing
  - ☒ Diskette with Sequence Listing in C.R.F.
  - ☒ Statement Accompanying Sequence Listings



09/869106

JC03 Rec'd PGT/PTC 22 JUN 2001

U.S. APPLICATION NO.

INTERNATIONAL APPLICATION NO.

ATTORNEY DOCKET NUMBER

Unassigned

PCT/FR99/03270

045636-5048

15. [X] The following fees are submitted:
- Basic National Fee (37 CFR 1.492(a)(1)-(5)):**
- Search Report has been prepared by the EPO or JPO.....\$860.00
- International preliminary examination fee paid to  
USPTO (37 CFR 1.482).....\$\*\*\*\*\*.00
- No international preliminary examination fee paid to  
USPTO (37 CFR 1.482) but international search fee  
paid to USPTO (37 CFR 1.445(a)(2)).....\$\*\*\*\*\*.00
- Neither international preliminary examination fee  
(37 CFR 1.482) nor international search fee  
(37 CFR 1.445(a)(2)) paid to USPTO.....\$\*\*\*\*\*.00
- International preliminary examination fee paid to USPTO  
(37 CFR 1.482) and all claims satisfied provisions  
of PCT Article 33(2)-(4).....\$\*\*\*\*\*.00

ENTER APPROPRIATE BASIC FEE AMOUNT = \$860.00

Surcharge of \$130.00 for furnishing the oath or declaration later than

[ ] 20 [ ] 30 months from the earliest claimed priority date

(37 CFR 1.492(e)).

Claims	Number Filed	Number Extra	Rate	
Total Claims	12 - 20 =	0	X \$18.00	\$0.00
Independent Claims	2 - 3 =	0	X \$80.00	\$0.00
Multiple dependent claim(s) (if applicable)			+\$270.00	\$0.00

TOTAL OF ABOVE CALCULATIONS = \$860.00

Reduction by 1/2 for filing by small entity, if applicable. Verified

Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28)

SUBTOTAL = \$0.00

Processing fee of \$130.00 for furnishing the English translation later

than [ ] 20 [ ] 30 months from the earliest claimed priority date

(37 CFR 1.492(f)).

TOTAL NATIONAL FEE = \$0.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The

assignment must be accompanied by an appropriate cover sheet

(37 CFR 3.28, 3.31).

\$40.00 per property

TOTAL FEES ENCLOSED = \$0.00

Amount to be

refunded

charged

\$

\$

\$

- a. [ ] A check in the amount of \$-0- to cover the above fees is enclosed.
- b. [X] Please charge my Deposit Account No. 50-0310 for \$860.00 to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. [X] **Except** for issue fees payable under 37 C.F.R. §1.18, the Commissioner is hereby authorized by this paper to charge any additional fees during the entire pendency of this application including fees due under 37 CFR §1.16 and §1.17 which may be required, or credit any overpayment to Deposit Account No. 50-0310.

Customer No. 009629

SEND ALL CORRESPONDENCE TO:

Morgan, Lewis & Bockius LLP

1800 M Street, N.W.

Washington, D.C. 20036

(202) 467-7000

*Lawrence J. Carroll*

Lawrence J. Carroll, Ph.D.

Reg. No. 40,940

Submitted: June 22, 2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: )  
 )  
Isabelle POQUET *et al.* )  
 )  
Application No.: ) Group Art Unit: Unassigned  
(based on PCT/FR99/03270) )  
 )  
Filed: Herewith ) Examiner: Unassigned  
 )  
For: *Gram-Positive Bacteria Deprived of HtrA* )  
*Protease Activity, and Their Uses* )

Commissioner for Patents and Trademarks  
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Dear Sir:

Prior to examination on the merits, please make the following changes to this application.

IN THE CLAIMS

Please cancel claims 1-10.

Please insert the following new claims 11-22.

--11. A method of producing a protein of interest, comprising:  
culturing a bacterial strain that expresses the protein, wherein the bacterial strain is a  
gram positive strain and wherein the bacterial strain does not express a functional HtrA  
protease.

12. A method according to claim 11, wherein the size of the genome of the bacterial  
strain is equal to or less than 3.2 Mb.

13. A method according to claim 11, wherein the bacterial strain is selected from a  
group consisting of *Streptococcaceae*, *Lactobacillaceae*, *Bacillaceae* of the  
*Staphylococcus* genus, *Bacillaceae* of the *Listeria* genus, and *Enterococcaceae* of the  
*Enterococcus* genus.

14. A method according to claim 11, wherein the bacterial strain is selected from a group consisting of *Lactococcus* spp., *Lactobacillus* spp., and *Streptococcus thermophilus*.
15. A method according to claim 11, wherein the bacterial strain does not express a functional PrtP protease.
16. A method according to claim 11, further comprising:  
introducing into the bacterial strain a nucleic acid having a sequence encoding a protein of interest operably linked to a promoter, wherein the nucleic acid is not integrated into a gene encoding the HtrA protease; and  
culturing the bacterial strain under conditions causing expression of the protein from the nucleic acid.
17. A Gram positive bacterial strain that does not express a functional HtrA protease wherein the strain has a genome that is equal to or less than 3.2 Mb in size, with the proviso that the bacterial strain is not a *Lactobacillus helveticus* strain having a *gusA* reporter gene inserted into a gene encoding the HtrA protease, and wherein the bacterial strain comprises an expression cassette having a sequence encoding a protein of interest operably linked to a promoter.
18. A bacterial strain according to claim 17, wherein the strain does not express a functional PrtP protease.
19. A method of producing a fermented product, comprising:  
culturing a bacterial strain according to claim 17 with a fermentation substrate under conditions suitable to produce a fermented product.
20. A method of producing a dietetic food, comprising:  
culturing a bacterial strain according to claim 17 with a substrate under conditions suitable to produce a dietetic food.
21. A method of producing a medicinal product, comprising:

culturing a bacterial strain according to claim 17 with a substrate under conditions suitable to produce a medicinal product.

22. A method according to claim 21, wherein the medicinal product is a vaccine.

### REMARKS

Applicants respectfully submit that no prohibited new matter has been introduced by this Preliminary Amendment and that claims 11-22 are drawn to the same invention as claims 1-10 of International Application PCT/FR99/03270. The changes to the claims represent changes in formalities so as to bring the claims into compliance with the rules of practice in the United States, such as "use" claims (original claims 7-10) have been re-written as method claims; standard claim terminology has been adopted i.e., "characterized in that" has been replaced by comprising (see original claims 1 and 6) and to adopt standard grammatical constructions (see all the original claims). The limitation with regard to genome size present in original claim 1 has been removed from claim 11 and placed in dependent claim 12. These changes do not narrow the scope of the claimed subject matter present and examined in the corresponding International Application.

Respectfully submitted,  
**MORGAN, LEWIS & BOCKIUS LLP**

Dated: June 27, 2001

By: 

Lawrence J. Carroll, Ph.D.  
Registration No. 40,940

**Customer No. 009629**  
**MORGAN, LEWIS & BOCKIUS LLP**  
1800 M Street, NW  
Washington, D.C. 20036  
Tel: 202-467-7000; Fax: 202-467-7258

9/18/84

09/869106

JC03 Rec'd PCT/PTC 22 JUN 2001  
PCT/FR99/03270

WO 00/39309

GRAM-POSITIVE BACTERIA DEPRIVED OF HtrA PROTEASE  
ACTIVITY AND THEIR USES

The invention relates to the production, in Gram-positive bacteria, of exported proteins.

5       The general term "exported proteins" denotes proteins which are transported across the cytoplasmic membrane. In the case of Gram-positive bacteria, this transport results in the secretion of the protein into the medium, or its association with the cell surface.

10       One of the main problems which arises during the production of exported proteins of interest by host bacteria lies in the degradation of these proteins during and/or after their exportation, at the cell envelope or at the cell surface. This degradation often  
15 leads to a decrease in the yield, and/or a modification of the structure and of the activity of the protein.

      The enzymes responsible for this degradation of exported proteins are bacterial proteases, themselves exported in the envelope; they are "housekeeping"  
20 proteases, one of the main functions of which is normally a role of degradation of abnormal or incorrectly folded exported proteins which accumulate in the medium or in the envelope, in particular under conditions of stress, and the role of which is also the  
25 recycling of exported proteins.

      Heterologous proteins, which are often incompletely recognized by the chaperone proteins involved in protein folding in the host bacterium, are particularly sensitive to attack by these proteases.

30       The oldest characterized exported housekeeping protease is the *E. coli* serine protease HtrA/DegP. It is a protease which as a periplasmic location, and which is expressed under the control of a promoter which is inducible at high temperature; Beckwith and  
35 Strauch (Proc. Natl. Acad. Sci. USA 85:1576-1580, 1988) have observed that it is involved in the proteolysis of proteins made from fusion between exported proteins of *E. coli* and the *PhoA* exportation reporter. They have proposed the inactivation of this protease in *E. coli*

09869106-100904

in order to limit the degradation of the heterologous exported proteins.

Mutant *E. coli* strains, in which the gene encoding the HtrA/DegP protease has been inactivated, have thus been obtained [Beckwith and Strauch, abovementioned publication, and PCT application WO 88/05821]; however, it has been noted that this inactivation results in a slowing down of the kinetics of degradation, but is not sufficient to abolish it because of the existence, in the envelope, of other proteases which degrade the exported proteins.

In *E. coli* several envelope housekeeping proteases, which carry out functions similar to those of HtrA/DegP, have been characterized: they are in particular the HhoA/DegQ and HhoB/DegS proteases, which are structurally homologous to HtrA/DegP, and proteases which are structurally different but functionally comparable (ApeA/proteaseI, OmpT, OmpP, Prc/Tsp, SppA/proteaseIV, PrtIII and SohB).

Studies relating to other bacteria have also made it possible to demonstrate the existence, in each species studied, of several exported housekeeping proteases. For example, a large number of bacterial species have several proteases of the HtrA family (Pallen and Wren, Mol. Microbiol. 19:209-21, 1997); three homologues of HtrA have been identified in *B. subtilis* (YyxA, YkdA and YvtB/Yirf), *Synechocystis* (HtrA, HhoA and HhoB), *Pseudomonas aeruginosa* and *Aquifex aeolicus*, two in *Hemophilus influenzae* (HtoA and HhoB), *Campylobacter jejuni*, *Brucella abortus* and *Yersinia enterocolitica*, and four in *Mycobacterium tuberculosis*. Various Gram-positive bacteria also have serine proteases considered to be related to the HtrA family on the basis of homology in the catalytic domain: EtA, EtB and V8/StsP of *S. aureus*, GseP of *Bacillus licheniformis* and Spro of *Mycobacterium paratuberculosis* (Koonin et al., Chap 117 in *Escherichia coli* and *Salmonella typhimurium*, 2203-17, 1997). Finally, exported proteases which are not

related to HtrA have also been demonstrated, for example in *B. subtilis* (Margot and Karamata, Microbiology, 142:3437-44, 1996; Stephenson and Harwood Appl. Environn. Microbiol. 64:2875-2881, 1998; Wu et al. J. Bacteriol. 173:4952-58, 1991).

It has therefore been proposed to combine mutations affecting several exported proteases in order to obtain an effective decrease in the degradation of heterologous exported proteins.

For example, an *E. coli* strain mutated in the *degP/htrA*, *ompT*, *prt* and *prc* genes (Meerman and Georgiou, Bio/technology 12:1107-10, 1994), and a *B. subtilis* strain deficient in the six extracellular proteases (Wu et al., 1991, abovementioned publication), have been constructed with this aim. However, the use of these strains does not make it possible to completely eliminate the proteolysis of the exported proteins. For example, in the case of the *B. subtilis* strain described by Wu et al., although the residual extracellular protease activity is negligible (< 1%), degradation of the heterologous exported proteins remains significant. In order to overcome this problem, that same team has carried out further modifications to this strain in order to make it overproduce various chaperones (Wu et al., J. Bacteriol. 180:2830-35, 1998). Furthermore, although the inactivation of the gene of one of these exported housekeeping proteases does not have any notable consequences for the bacterium, the accumulation of mutations may affect strain viability; Meerman and Georgiou (1994, abovementioned publication) thus observe a decrease in growth rate which can range up to 50%.

In lactic acid bacteria, only a few exported proteases have been studied; the most well characterized at the present time is the protease named PrtP (Kok, FEMS Microbiol. Reviews 87:15-42, 1990), which is located at the cell surface, where it is anchored to the peptidoglycan. This protease is present



in many lactic acid bacteria, in particular *Lactococcus lactis*, and is located on a plasmid. It contributes to the nitrogen-based nutrition of bacteria by degrading milk caseins. Other surface proteases have been purified from two species of lactic acid bacteria, *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Lactobacillus helveticus*, but their function has not been determined (Stefanitsi et al., FEMS Microbiol. Lett. 128:53-8, 1995; Stefanitsi and Garel, Lett. Appl. Microbiol. 24:180-84, 1997; Yamamoto, et al., J. Biochem. 114:740-45, 1993). A stress-induced gene encoding a protein which is highly homologous to the proteases of the HtrA family has recently been revealed in *Lactobacillus helveticus* (Smeds et al., J. Bacteriol. 180:6148-53, 1998). It has been observed that this gene is necessary for survival at high temperature; a mutant *Lactobacillus helveticus* strain in which the *htrA* gene has been inactivated by insertion of a reporter gene (*gusA*, encoding  $\beta$ -glucuronidase) under the control of the *htrA* promoter, was constructed. The study of the expression of the *gusA* gene in this mutant made it possible to demonstrate induction of the transcription of this gene under the same conditions as that of the *htrA* gene in the wild-type strains; on the other hand, no  $\beta$ -glucuronidase activity was observed.

In previous investigations directed towards studying exported proteins of *Lactococcus lactis* by studying proteins fused with the  $\Delta_{SP}Nuc$  exportation reporter (Poquet et al., J. Bacteriol. 180:1904-12, 1998), the team of inventors has observed significant extracellular proteolysis even though the experiments were carried out in an *L. lactis* subsp. *cremoris* strain free of any plasmid and therefore, in particular, of that which carries *prtP*.

The inventors undertook to investigate extracellular proteases responsible for this proteolysis.

They have thus discovered, in *L. lactis*, the existence of a gene of the *htrA* family.

This gene, detected in the genome of the IL1403 strain of *L. lactis* subsp. *lactis*, encodes a 408 amino acid protein, hereinafter named HtrA<sub>L1</sub>, the nucleotide sequence and the amino acid sequence of which are represented on figure 1, and appear in the attached sequence listing (SEQ ID NO: 1). This protein is very homologous to *E. coli* HtrA, and to various other known members of the HtrA family, as shown in table I below, which illustrates the percentages of identity and of similarity between HtrA<sub>L1</sub> and various proteins of the HtrA family:

TABLE I

Protein	Organism	% identity	% similarity
HtrA/DepP/Do protease	<i>E. coli</i>	31.5	38.2
HhoA/DegQ	<i>E. coli</i>	34.0	40.8
HhoB/DegS	<i>E. coli</i>	29.9	37.3
HtrA	<i>S. typhimurium</i>	32.4	39.1
HtoA	<i>H. influenzae</i>	31.9	39.2
HhoB/DegS	<i>H. influenzae</i>	31.2	40.0
spHtrA	<i>S. pneumoniae</i>	55.6	62.0
HtrA	<i>Lb. helveticus</i>	46.9	54.1
YyxA	<i>B. subtilis</i>	43.5	52.0
YkdA	<i>B. subtilis</i>	42.5	49.4

The HtrA protein of the IL1403 strain of *L. lactis* subsp. *lactis* has the three amino acids Ser, His and Asp, which define the catalytic site characteristic of serine proteases related to trypsin, among which is the HtrA family; in addition, it has, around these three amino acids, the following three motifs: DAYVVTNYH<sub>127</sub>VI, D<sub>157</sub>LAVLKIS, and GNS<sub>239</sub>GGALINIEGQVIGIT, which correspond to the consensus regions defined by Pallen and Wren (Mol. Microbiol. 19:209-21, 1997) for the catalytic domain of the HtrA proteases: -GY--TN-HV-, D-AV---- and GNSGG-L-N-G--IGIN.

At its N-terminal end, it has a hydrophobic amino acid sequence L<sub>10</sub>LTGVVGGAIALGGSAI<sub>26</sub> corresponding

09669105-100901

to a putative transmembrane segment. The HtrA<sub>L1</sub> protein of *L. lactis* subsp. *lactis* is therefore thought to be an integral protein of the cytoplasmic membrane. According to the "positive inside" rule concerning the topology of these proteins (Von Heijne, Nature, 341:456-8, 1989), its topology corresponds to the "C-out" type, i.e. its C-terminal portion, which comprises in particular its catalytic site, would be exposed to the outside of the plasma membrane. Like the HtrA protease of *E. coli*, *L. lactis* subsp. *lactis* HtrA<sub>L1</sub> therefore appears to be an envelope protease which can degrade exported proteins. The amino acids of the catalytic domain and of the transmembrane domain are framed on figure 1.

The inventors have inactivated this gene by mutation; at optimum temperature (30°C), the mutant *L. lactis* subsp. *lactis* strain thus obtained is viable and grows normally; on the other hand, its growth and viability are affected at higher temperatures (from 37°C), both on plates and in liquid medium.

In addition, the inventors have studied the effect of this mutation on the exportation of various fusion proteins, and have noted that the inactivation of the HtrA<sub>L1</sub> protease in *L. lactis* is sufficient to completely abolish the degradation of the exported proteins; this effect is surprising given the residual proteolysis observed previously in other bacteria after inactivation on proteases of the HtrA family.

A subject of the present invention is a process for producing a protein of interest, characterized in that it comprises culturing a bacterial strain which expresses said protein of interest, and which can be obtained from a Gram-positive bacterium, the size of the genome of which bacterium is at most equal to 3.2 Mb, preferably at most equal to 3 Mb, and advantageously at most equal to 2.5 Mb, by mutation which inactivates the HtrA surface protease of said bacterium;

and producing said protein of interest exported by said strain.

According to a preferred embodiment of the present invention, the starting Gram-positive bacterium is chosen from bacteria of the group consisting of the *Streptococcaceae*, and *Lactobacillaceae*. Advantageously, it is chosen from lactococci.

It may be also be chosen from bacteria belonging to the group consisting of the *Bacillaceae*, for example to the *Listeria* genus, and the *Enterococcaceae*, in particular of the *Enterococcus* genus.

Advantageously, said bacterial strain may also comprise one or more other modifications of its genome, directed toward improving the production and/or secretion of proteins expressed in said bacterium, and/or toward avoiding their degradation. Depending on the type of protein intended to be produced, it is possible, for example, to use a bacterial strain in which the PrtP protease activity has been inactivated, and/or a bacterial strain which overproduces a protein allowing the stabilization of exported proteins, such as the Nlp4 protein of *Lactococcus lactis*, or a homologue thereof (Poquet et al. 1998, abovementioned publication).

A subject of the present invention is also any bacterial strain which can be obtained from a Gram-positive bacterium, the size of the genome of which bacterium is at most equal to 3.2 Mb, as defined above, by mutation which inactivates the HtrA surface protease of said bacterium, and which also comprises at least one cassette for expressing a gene of interest, with the exception of a *Lactobacillus helveticus* strain comprising a single expression cassette consisting of the sequence encoding the *gusA* reporter gene inserted into the *htrA* gene of said strain, under the transcriptional control of the promoter of said gene.

The term "expression cassette" is intended to mean any recombinant DNA construct comprising a gene of

09869406-100904  
T06500T-90T69860

5 host bacterium under consideration.

10 *E. coli* which are sufficient for it to be included in  
the same family, i.e.:

- a catalytic site formed by the three amino acids Ser, His and Asp;

15       - the presence, around this catalytic site, of  
the consensus regions: -GY--TN-HV-, D-AV---- and  
GNSGG-L-N-G-IGIN;

- an exportation signal enabling the protease to be transported to the cell surface of the bacterium, (it may, for example, be a signal peptide, a transmembrane domain, a signal for anchorage to the wall, etc.).

In order to implement the present invention, mutant bacteria lacking HtrA activity can be produced by carrying out one or more mutations, in particular in the sequence encoding the HtrA protease and/or in the regulatory sequences allowing the expression of the *htrA* gene, so as to prevent the expression of a functional HtrA protease. These mutations can be carried out conventionally, by deletion, insertion or replacement of at least one nucleotide or one nucleotide sequence in the *htrA* gene; they can result either in the absence of production of HtrA, or in the production of an HtrA protease in which at least one amino acid required for activity has been deleted or replaced.

The suitable mutagenesis techniques are known per se; advantageously, use will be made of site-directed mutagenesis techniques, since the data available on the proteases of the HtrA family make it

possible, even though more precise information on the specific sequence of the gene whose inactivation is desired is not available, to target the mutation(s) on conserved domains which are required for activity (for  
5 example the catalytic domain).

The present invention can be implemented in many domains.

Firstly, it can be used in the domain of the production of proteins of interest (for example  
10 enzymes, human proteins, etc.) by genetic engineering, using cultures of bacteria transformed with a gene of interest. In this domain, the present invention makes it possible to improve the yield of exported proteins (and in particular secreted proteins), and to avoid  
15 their contamination with inactive proteolytic products: this makes it possible to purify them easily and less expensively.

For this application, use will preferably be made of the mutant strains produced from nonpathogenic  
20 bacteria, such as *Lactococcus* spp. or *Lactobacillus* spp., and also food streptococci, *Streptococcus thermophilus*.

The mutant strains produced from bacteria conventionally used in the agro-foods industry, such as  
25 lactic acid bacteria (in particular lactococci, lactobacilli and thermophilic streptococci), can advantageously be used in this domain. For example, they can be used in the composition of ferments, in order to produce heterologous proteins making it  
30 possible to improve the quality of the finished fermented product; thus, the exportation of foreign enzymes produced by a mutant *L. lactis* strain in accordance with the invention, within cheeses fermented with *L. lactis*, may improve their maturing and their  
35 organoleptic qualities.

These mutant strains can also be used for producing dietetic products or medicinal products. In this domain, mutant strains in accordance with the invention can, for example, be used in order to

09669106-100901

express, prior to the ingestion of the product and/or after its ingestion, proteins with a prophylactic or therapeutic effect, such as enzymes (for facilitating digestion, for example), proteins for stimulating the immune system, immunization antigens, etc. In most cases, for use in this domain, and in order to guarantee maximum innocuity, mutant strains produced from nonpathogenic bacteria and, advantageously, from bacteria conventionally used for food will be preferred. However, in the context of uses for immunization, mutant strains produced from pathogenic bacteria (in particular streptococci, staphylococci, enterococci or listeria), and preferably from variants of these bacteria already carrying one or more mutations which attenuate their pathogenic power, can be used; the inactivation of the HtrA protein, in limiting the capacities of survival of these bacteria under conditions of stress, may contribute to attenuating their virulence, as previously observed in the case of certain Gram-negative bacteria.

In the context of certain applications, in which the host bacterium must be viable and capable of producing proteins at temperatures of about 35 to 40°C, for example the production, in a fermentor, of certain proteins, or the production, after ingestion, in the digestive tract of humans or animals, of proteins with therapeutic or prophylactic activity, mutant strains produced from thermophilic bacteria, such as *Streptococcus thermophilus*, will advantageously be used.

The present invention will be more clearly understood with the aid of the continuation of the description which follows, which refers to nonlimiting examples illustrating the production of *L. lactis* mutants in which the HtrA surface protease is inactive, and the properties of these mutants.

**EXAMPLE 1: INACTIVATION OF THE *htrA* gene of *L. lactis***

*htrA* gene, carried by the chromosome of the IL1403 strain (Chopin et al. Plasmid, 11, 260-263,

05869106-100901  
T0600T-9016880

1984) of *L. lactis* subsp. *lactis*, was inactivated by integration of a suicide plasmid carrying a 665 bp internal fragment of the gene (FA).

As a positive control for integration, a suicide plasmid carrying a 902 bp fragment truncated in the 3' region (GA), the integration of which onto the chromosome restores a wild-type copy of the gene, was used.

These fragments were obtained beforehand by PCR amplification from the genomic DNA of the IL1403 strain of *L. lactis* subsp. *lactis*, using the pairs of primers F/A and G/A:

- F[5'-GGAGCCA(G/T)(A/C/T)GC(A/G/C/T)(C/T)T(A/G/T)GG-3']  
located downstream of the ATG initiation codon  
- G[5'-GTTTCCACTTTTCTGTGG-3']  
located upstream of the *htrA* promoter  
- A[5'-TT(A/T)CC(A/T)GG(A/G)TT(A/G/T)AT(A/G/C/T)GC-3']  
located upstream of the serine codon of the catalytic site.

The positioning of the F, G and A primers is indicated on figure 1.

The amplification was carried out under the following conditions:

- reaction mixture: 0.2 mM of each dNTP, 5  $\mu$ M of each oligonucleotide, approximately 500 ng of chromosomal DNA, 2 mM of MgCl<sub>2</sub> and 1.25 units of Taq-DNA-pol (Boehringer Mannheim), in the Taq buffer provided by the manufacturer;

- temperature conditions: 5 min 94°C, 30 cycles (30 sec at 94°C, 30 sec at 46°C, and 30 sec at 72°C), and 4°C.

The amplified fragments were ligated to the linear pGEM<sup>T</sup> plasmid (Promega). After transformation of *E. coli* TG1 with the ligation products, the clones which are resistant to ampicillin and lack  $\beta$ -galactosidase activity are selected. The plasmids obtained, bearing the FA and GA fragments, respectively, are named pES1.1 and pES2.1.



The FA and GA inserts were subcloned into a suicide vector carrying a chloramphenicol resistance gene. Since this vector is incapable of replicating alone in the absence of the RepA protein which is required for initiating its replication, co-integrants were created by ligation between each of the pES1.1 and pES2.1 plasmids and the suicide vector, linearized beforehand.

After transformation of the *E. coli* TG1 strain, and selection of the chloramphenicol-resistant clones, the pGEM<sup>T</sup> portion of the co-integrants was deleted and the vectors were re-circularized. The plasmids obtained are multiplied in the TG1 repA<sup>+</sup> strain of *E. coli*; after selection of the chloramphenicol-resistant clones, the suicide plasmids named pVS6.1 and pVS7.4 are obtained.

pVS6.1 contains the FA fragment, and pVS7.4 contains the GA fragment, of the *htrA<sub>L1</sub>* gene of the IL1403 strain of *L. lactis* subsp. *lactis*.

These plasmids were used to transform the IL1403 strain of *L. lactis* subsp. *lactis*; the clones which had integrated these plasmids at the *htrA* locus on the chromosome were selected in the presence of chloramphenicol.

In both cases, several independent chloramphenicol-resistant clones were obtained. Five clones of each class termed A to E in the case of the integration of pVS6.1, and 17 to 22 in the case of the integration of pVS7.4, were chosen for analysis.

For each of these clones, the integration at the *htrA* locus was confirmed by Southern transfer.

Two clones, A and 17, were chosen for the following analyses; they constitute the two prototypes of the mutant strains, which hereinafter will be named:

- *htrA* (null mutation of the *htrA<sub>L1</sub>* gene, Cm<sup>R</sup>); this strain does not express any active HtrA protease;
- *htrA<sup>+</sup>/htrA* (wild-type copy + truncated copy of the *htrA<sub>L1</sub>* gene, Cm<sup>R</sup>); this strain expresses an active HtrA<sub>L1</sub> protease.

**EXAMPLE 2: ROLE OF THE *htrA*<sub>LI</sub> GENE OF *L. lactis* IN SURVIVAL AT HIGH TEMPERATURE**

The two strains *htrA* and *htrA*<sup>+</sup>/*htrA* are cultured, in liquid culture, under the conventional conditions for growth of *L. lactis*, i.e. at 30°C and in the presence of oxygen, but without stirring, and in the presence of chloramphenicol.

The behavior of the *htrA* strain of *L. lactis* subsp. *lactis* at 30°C and at 37°C was studied using the *htrA*<sup>+</sup>/*htrA* strain and also the IL403 parent-strain (cultured in the absence of chloramphenicol) as control.

The bacteria were cultured overnight at room temperature, in an M17 medium containing 1% of glucose (+ 2.5 µg/ml of chloramphenicol for both the *htrA* strain and the *htrA*<sup>+</sup>/*htrA* strain). The cultures were diluted 100-fold in the morning, in the same medium, and divided into two batches placed in semi-anaerobiosis at 30°C or at 37°C. The growth was monitored by measuring the OD<sub>600</sub>.

The results are illustrated in figure 2.

At 30°C (fig. 2A), it is noted that the *htrA*<sup>+</sup>/*htrA* strain (■), the *htrA* strain (◆), and the wild-type IL1403 strain (▲) have very close generation times: 65 min for the wild-type strain, 70 min for *htrA*<sup>+</sup>/*htrA* and 75 min for *htrA*; finally, for the 3 cultures, the OD<sub>600</sub> values corresponding to the stationary phase are very comparable (OD<sub>600</sub> = 2.1 to 2.2).

These results indicate that there is no significant difference in growth between these three strains at 30°C.

At 37°C (fig. 2B), the *htrA*<sup>+</sup>/*htrA* strain (■) has a generation time of 100 min and the OD<sub>600</sub> of the stationary phase is less than at 30°C (OD<sub>600</sub> = 1.25). Less growth at 37°C than at 30°C is also observed for the wild-type IL1403 strain (▲); the generation time is 65 min, but the OD<sub>600</sub> of the stationary phase is less than at 30°C (OD<sub>600</sub> = 1.9). In the case of the *htrA*

strain (◆), the growth is very slight, or even zero, and the OD<sub>600</sub> does not exceed 0.1, even after culturing for 7 h.

5 It emerges from these results that the *htrA* strain of *L. lactis* subsp. *lactis* is heat-sensitive and that the *htrA* mutation is lethal at 37°C.

**EXAMPLE 3: ROLE OF THE *htrA<sub>LI</sub>* GENE OF *L. LACTIS* IN SURFACE PROTEOLYSIS**

10 The effect of the *htrA<sub>LI</sub>* mutation on the stability of five exported proteins was tested. These proteins are:

15 i) a heterologous protein, the secreted nuclease of *S. aureus*, Nuc; this protein is expressed by the plasmid pNuc3 (Le Loir et al., J. Bacteriol. 176:5135-5139, 1994; Le Loir et al., J. Bacteriol. 180:1895-903, 1998);

20 ii) three hybrid proteins (Usp-Δ<sub>Sp</sub>Nuc, Nlp4-Δ<sub>Sp</sub>Nuc and Exp5-Δ<sub>Sp</sub>Nuc) resulting from the fusion between the Δ<sub>Sp</sub>Nuc reporter and fragments of exported proteins of *L. lactis*: the secreted protein Usp45 (Van Asseldonk et al., Gene 95:155-60, 1990), the lipoprotein Nlp4 and the protein Exp5 (which is, itself, a protein made from fusion between an exported protein and a cytoplasmic protein); these proteins, and  
25 also the plasmids pVE8009, pVE8024 and pVE8021 which express them, respectively, are described by Poquet et al. (1998, abovementioned publication);

30 iii) a naturally exported protein of *L. lactis*, AcmA.

35 In the wild-type MG1363 strain of *L. lactis* subsp. *cremoris*, Usp-Δ<sub>Sp</sub>Nuc is secreted and Nlp4-Δ<sub>Sp</sub>Nuc is associated with the cells; for these two proteins, various degradation products, among which the NucA peptide originating from the Δ<sub>Sp</sub>Nuc portion of the fusion, are detected in the medium, along with the mature form; with regard to the Exp5-Δ<sub>Sp</sub>Nuc tripartite fusion, it is very unstable and the mature form is not detected in the medium, only the degradation products, including the NucA peptide. The mature form, and also

the degradation products of these three hybrid proteins, can be detected using anti-NucA antibodies.

The naturally exported protein of *L. lactis* chosen is the bacteriolysin AcmA (Buist et al., J. Bacteriol. 177:1554-1563, 1995). This protein, which degrades peptidoglycan, is both secreted and associated with the surface, probably by affinity with its substrate. It provides, both in the MG1363 strain of *L. lactis* subsp. *cremoris* and the IL1403 strain of *L. lactis* subsp. *lactis*, proteolysis products which are active and therefore detectable, like the intact protein, by zymogram.

The strains transformed with the plasmids expressing these various proteins are cultured at 30°C for several hours, at least up to the middle of the exponential phase or up to the start of the stationary phase.

For each plasmid, cultures of the three strains IL1403, *hrtA* and *htrA*<sup>+</sup>/*htrA*, which had reached comparable OD<sub>600</sub> values, were used to extract protein samples: a) from the total culture, b) from the cells and c) from the medium, according to the protocol described by Poquet et al. (1998, abovementioned publication).

These samples are subjected to electrophoresis (SDS-PAGE) on denaturing gel.

In order to detect the Nuc, USP- $\Delta_{SpNuc}$ , Nlp4- $\Delta_{SpNuc}$  and Exp5- $\Delta_{SpNuc}$  proteins and their degradation products, the proteins are transferred onto a membrane, followed by immunological revelation using anti-NucA antibodies, which are detected using a protein G/peroxidase conjugate (BIO-RAD) and a chemiluminescence kit (Dupont-Nen).

Acma is detected by zymogram (Buist et al., 1995, abovementioned publication): micrococci, in which the wall is sensitive to Acma, are included in the electrophoresis gel at the concentration of 0.2%, which makes it opaque; after electrophoresis, the gel is treated at 37°C overnight in a buffer containing 50 mM

of Tris/HCl at pH 7 and 0.1% of Triton X100, which allows lysis of the micrococci by AcmA or its active proteolytic products. The gel is then colored with methylene blue at 0.1% in 0.01% KOH: the bands  
5 corresponding to the AcmA activity appear as transparent hydrolysis halos on a blue background.

For each protein, the degradation profiles in the IL1403, *htrA* and *htrA<sup>+</sup>/htrA* strains were compared by observing the protein content accumulated during  
10 culturing for several hours.

Figures 3 to 6 show, respectively, the results of immunological detection for the Nuc, Usp- $\Delta_{Sp}Nuc$ , Nlp4- $\Delta_{Sp}Nuc$  and Exp5- $\Delta_{Sp}Nuc$  proteins. For the Nuc (fig. 3) and Usp- $\Delta_{Sp}Nuc$  (fig. 4) proteins, [lacuna]

15 Fig. 7 represents a zymogram of the bacteriolysin activity of AcmA; the detection was carried out on the total culture (T), the cells alone (C) or the medium (M).

In the IL1403 strain:

20 For the secreted proteins Nuc and Usp- $\Delta_{Sp}Nuc$  (fig. 3 and 4: first three wells), and for the lipoprotein Nlp4- $\Delta_{Sp}Nuc$  (fig. 5: first well), a three-band profile is detected, as previously observed in the MG1363 strain (Le Loir et al., 1994; Poquet et al.,  
25 1998, abovementioned publications):

a) the band with the highest molecular weight is the precursor from which the signal peptide has not been cleaved, which is confirmed by its presence exclusively in the cells (fig. 3 and 4);

30 b) the intermediate band is the mature form after cleavage of the signal peptide, and, in the case of the secreted proteins Nuc and Usp- $\Delta_{Sp}Nuc$  (fig. 3 and 4), it is present exclusively in the medium;

c) the band with the lowest molecular weight is  
35 the NucA peptide which practically comigrates with the commercial NucA form purified from *S. aureus* (the slight difference in migration being due to the different cleavage specificities in *S. aureus* and

*L. lactis*), and which is both released into the medium and associated with the cells.

For the Exp5- $\Delta_{SpNuc}$  protein (fig. 6: first well), two forms are detected only with great difficulty, one having a high molecular weight and one having a low molecular weight, NucA, which practically comigrates with the purified commercial form; there is, therefore, practically total proteolysis in IL1403.

For the AcmA protein (fig. 7: the first three wells), a four-band profile, as previously observed in the MG1363 strain (Buist et al., 1995, abovementioned publication), is detected:

a) the band with the highest molecular weight is the precursor from which the signal peptide has not been cleaved, which is present exclusively in the cells;

b) the band with a slightly lower molecular weight is the mature form after cleavage of the signal peptide, which is both secreted into the medium and associated with the surface of the cells by affinity for its substrate;

c and d) the two bands of lower molecular weight are active proteolytic products, both secreted into the medium and associated with the surface of the cells by affinity for their substrate.

In the *htrA*<sup>+</sup>/*htrA* strain:

(Fig. 3 and 4: last three wells, fig. 5 and 6: last well, and fig. 7: last three wells). The profiles observed are absolutely identical to those observed in the wild-type strain. The *htrA*<sup>+</sup>/*htrA* strain therefore exhibits a wild-type proteolytic phenotype which is explained by the wild-type copy of the *htrA*<sub>L1</sub> gene which it possesses.

In the *htrA* strain:

(Fig. 3 and 4: three central wells, fig. 5 and 6: central well, and fig. 7: three central wells).

In all cases, none of the proteolytic products are detected; simultaneously, the amount of mature

protein (or of high molecular weight protein in the case of Exp5- $\Delta_{SpNuc}$ ) increases.

These results show that the product of the *htrA<sub>L1</sub>* gene is clearly responsible for the degradation  
5 of the secreted proteins, and that its inactivation leads to the complete abolition of this degradation.

09869106-100901  
"0600T" 90159860

## CLAIMS

1. A process for producing a protein of interest, characterized in that it comprises:

- culturing a bacterial strain which expresses  
5 said protein of interest, and which can be obtained from a Gram-positive bacterium, the size of the genome of which bacterium is at most equal to 3.2 Mb, by mutation which inactivates the HtrA surface protease of said bacterium;
- 10 - producing said protein of interest exported by said strain.

2. The process as claimed in claim 1, characterized in that the starting Gram-positive bacterium is chosen from the *Streptococcaceae*,  
15 *Lactobacillaceae*, *Bacillaceae* of the *Staphylococcus* and *Listeria* genera, and *Enterococcaceae* of the *Enterococcus* genus.

3. The process as claimed in claim 2, characterized in that the starting Gram-positive  
20 bacterium is chosen from the group consisting of *Lactococcus* spp., *Lactobacillus* spp., and *Streptococcus thermophilus*.

4. The process as claimed in any one of claims 1 to 3, characterized in that the bacterial strain used  
25 also lacks PrtP protease activity.

5. A bacterial strain which can be obtained from a Gram-positive bacterium, the size of the genome of which bacterium is at most equal to 3.2 Mb, as defined in any one of claims 1 to 3, by mutation which  
30 inactivates the HtrA surface protease of said bacterium, and which also comprises at least one cassette for expressing a gene of interest, with the exception of a *Lactobacillus helveticus* strain comprising a single expression cassette consisting of  
35 the sequence encoding the *gusA* reporter gene inserted into the *htrA* gene of said strain, under the transcriptional control of the promoter of said gene.

09869106-100001



6. The bacterial strain as claimed in claim 5, characterized in that it also lacks PrtP protease activity.

7. The use of a bacterial strain as defined in any  
5 one of claims 1 to 4, for preparing a fermented product.

8. The use of a bacterial strain as defined in any one of claims 1 to 4, for preparing a dietetic food.

9. The use of a bacterial strain as defined in any  
10 one of claims 1 to 4, for preparing a medicinal product.

10. The use as claimed in claim 9, characterized in that said medicinal product is a vaccine.

09669106-100901

1/9




FIG. 1

```

661  agtcaaaagatttcgtggttggttatgatgaatatcacagacatccgtgctgtcttcaaaaatcag 720
      V K D S V V G Y D E Y T D L A V L K I S
                                catalytic domaine

721  ttcctgaacatgttcaaaagatgtgtggagacatccgtgtgattcttagtaaaattcaacaatttggtga 780
      S E H V K C V A T F A D S S K L T I G E

781  accttgccatttcgctgttggttacccttttaggttagtcaattcttgcaaacacccgcaattgaaagg 840
      P A I A V G S P L G S Q F A N T A T E G

841  aatcttatcttgcaacaaagctgtcaagtgaattcgatcccaagaaaaatggttcaaaataactaa 900
      I L S A T S R Q V T L T Q E N G Q T T N

                                primer A
901  tatcaaatgcaatttcaaacagatgcttgccatttaacccctggtaactcttgaggagggcctttgat 960
      I N A I Q T D A A I N P G N S G G A L I
                                catalytic domaine

961  taatatctgaaggacaaggttatctggaattactcaaaagtataaaattacaacaactgaagatgg 1020
      N I E G Q V I G I T Q S K I T T T E D G

1021  ttcctactcttgctgaaggctttaggattttcgatttccttcaatgatgtctgtaaacatcat 1080
      S T S V E G L G F A I P S N D V V N I I

1081  caataaaacttgaaagatgatggttaagatttcacagccctgtctttaggtatccgaaatgggtga 1140
      N K L E D D G K I S R P A L G I R M V D

1141  cctttcacaaattatcaacaaaatgacagttctcaattgaaattactaagcagtgtaaacagg 1200
      L S Q L S T N D S S Q L K L L S S V T G

1201  tgggggttggtgttttaattctgttcaaatctgggacttcctgtgtgctctcagctggtttgaaagg 1260
      G V V V V Y S V Q S C L P A A S A G L K A

1261  tggagatgttaattacaaaaggttgggtgacacagcagtaaacctcttcaaacagacttgcaaaag 1320
      G D V I T K V G D T A V T S S T D L Q S

```

3/9

1321 ----- 1380  
tgctctttactccacacaatatcaatgatacagtaaaagttaattattatcgtgatggtaa  
A L Y S H N I N D T V K V T Y Y R D G K  
1381 ----- 1440  
atcaaatcacagcagagatgttaaaactttttaaatacaaccagtgacttagaaacaagcagtc  
S N T A D V K L S K S T S D L E T S S P  
1441 ----- 1500  
atctctctcttaattaaacaacttaataatttaataaaagtctcttgtaaatagaaggcttt  
S S S N  
1501 ----- 1560  
tttcatactaaagtctgaaatcttttaaaaaataataatctccatctctctctctctgatt  
1561 ----- 1620  
tatggtaaaataaagttaagcagaaaaattttactttactttaagaaagccgaacaaattctg  
1621 ----- 1680  
agtcatcaggaattctggctgtgaaatgaacacatcaacaacggcctcttgatttaacgggc  
1681 ----- 1740  
attgacctggacaaaaaatcttgaggatgattacgatatctctccatttaaaatacttatggc


FIG. 1 (continued)

WO 00/39309

PCT/FR99/03270

4 / 9

A



B




FIG. 2

FIG. 3

Nuc




FIG. 3

Usp-Δ<sub>sp</sub>Nuc




FIG. 1

WO 00/39309

PCT/FR99/03270

10

Nlp4-Δ<sub>SP</sub>Nuc

precursor  
mature form

NucA  
hira/hira  
hira  
IL1403 VIT  
NucA

FIG. 5

09869106-100904



intact form → 

Nir A


IL1A03:WT  
hira  
hira+hira  
Nucca

FIG. 6



FIG. 7

# COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY



U.S. DEPARTMENT OF COMMERCE  
Patent and Trademark Office

ATTORNEY DOCKET NO. :

As a below named inventor, I hereby declare that :

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled :

## GRAM-POSITIVE BACTERIA DEPRIVED OF HtrA PROTEASIC ACTIVITY AND THEIR USES

the specification of which :

is attached hereto ; or

was filed as United States application Serial No. \_\_\_\_\_ on \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable) ; or

was filed as a PCT international application Number PCT/FR99/03270 on December 23, 1999 and was amended under PCT article 19 on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose the U.S. Patent and Trademark Office information which is material to the patentability of claims presented in this application in accordance with Title 37, Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate or §365(a) of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed :

### PRIOR FOREIGN APPLICATION(S) :

COUNTRY (if PCT, indicate PCT)	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED
FRANCE	98/16462	24 December 1998	X

Combined Declaration For Patent Application and Power of Attorney – (Continued)  
(includes Reference to PCT International Applications)

ATTORNEY DOCKET NO. :

I hereby claim the benefits under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

U.S. PROVISIONAL APPLICATIONS

U.S. PROVISIONAL APPLICATION NO.

U.S. FILING DATE

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or §365 (c) of any PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose to the U.S Patent and Trademark Office all information known to me to be material to the patentability of claims represented in this application in accordance with Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application(s) and the national or PCT international filing date of this application :

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT :

U.S. APPLICATIONS

STATUS (Check one)

U.S. APPLICATION NO.

U.S. FILING DATE

PATENTED

PENDING

ABANDONED

**POWER OF ATTORNEY:** As a named inventor, I hereby appoint the registered practitioners of Morgan, Lewis & Bockius LLP included in the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct all correspondence be addressed to that Customer Number.

**Customer Number : 009629**

Direct Telephone Calls To :  
(name and telephone number)

Combined Declaration For Patent Application and Power of Attorney – (Continued)  
(includes Reference to PCT International Applications)

ATTORNEY DOCKET NO. :

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

FULL NAME OF SOLE OR FIRST INVENTOR	<b>POQUET Isabelle</b>	
RESIDENCE & CITIZENSHIP	<b>PARIS, FRENCH</b> <i>FRX</i>	COUNTRY OF CITIZENSHIP <b>FRANCE</b>
POST OFFICE ADDRESS	56-62, Rue de Vouillé 75015 PARIS FRANCE	
FIRST OR SOLE INVENTOR'S SIGNATURE	<i>Isabelle Poquet</i>	DATE <b>28.06.01</b>
FULL NAME OF SECOND INVENTOR	<b>GRUSS Alexandra</b>	
RESIDENCE & CITIZENSHIP	<b>ORSAY, AMERICAN</b> <i>FRX</i>	COUNTRY OF CITIZENSHIP <b>THE UNITED-STATES</b>
POST OFFICE ADDRESS	25, Rue Louis Scocard 91400 ORSAY FRANCE	
SECOND INVENTOR'S SIGNATURE	<i>Alexandra Gruss</i>	DATE <b>05-07-01</b>
FULL NAME OF THIRD INVENTOR	<b>BOLOTINE Alexandre</b>	
RESIDENCE & CITIZENSHIP	<b>NANCY, RUSSIAN</b> <i>FRX</i>	COUNTRY OF CITIZENSHIP <b>FEDERATION OF RUSSIA</b>
POST OFFICE ADDRESS	5, Rue Maréchal Galliéni 54000 NANCY FRANCE	
THIRD INVENTOR'S SIGNATURE	<i>Alexandre Bolotine</i>	DATE <b>06.07.2001</b>
FULL NAME OF FOURTH INVENTOR	<b>SOROKINE Alexei</b>	
RESIDENCE & CITIZENSHIP	<b>GIF-SUR-YVETTE, RUSSIAN</b> <i>FRX</i>	COUNTRY OF CITIZENSHIP <b>FEDERATION OF RUSSIA</b>
POST OFFICE ADDRESS	8, Résidence les Quinconces 91190 GIF-SUR-YVETTE FRANCE	
FOURTH INVENTOR'S SIGNATURE	<i>Alexei Sorokine</i>	DATE <b>06.07.2001</b>

Listing of Inventors Continued on attached page(s) [ ] Yes [ X ] No

09/869106

JC03 Rec'd PCT/PTC 22 JUN 2001

1

## SEQUENCE LISTING

<110> INSTITUT NATIONAL DE LA RECHERCHE AGRONOMIQUE (INRA)  
 POQUET, Isabelle  
 GRUSS, Alexandra  
 BOLOTINE, Alexandre  
 SOROKINE, Alexei

<120> GRAM-POSITIVE BACTERIA DEPRIVED OF HtrA PROTEASIC  
 ACTIVITY, AND THEIR USES

<130> MJPcb539/89

<140>

<141>

<150> FR9816462

<151> 1998-12-24

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 1740

<212> DNA

<213> Lactococcus lactis

<220>

<221> CDS

<222> (230)..(1453)

<400> 1

aaacaagatg aaaacatgat ttatcaacat ttttttactt ttttccactt ttctgtggaa 60  
 aactttatta aaatatccac ttatcctcat taatttttag attatccaca aaaatgtgga 120  
 gaaactatat tagtttgatt tttgttacta ttaaggtatt attaagtgag agtagatata 180  
 attacatcat agaaatgcta caaagattaa taattgaaag gaattatgt atg gca aaa 238  
 Met Ala Lys  
 1

gct aat ata gga aaa ttg cta tta aca ggt gtc gtg ggc gga gcc atc 286  
 Ala Asn Ile Gly Lys Leu Leu Leu Thr Gly Val Val Gly Gly Ala Ile  
 5 10 15

gca ctt gga gga agt gca atc tat caa agc act aca aat caa tcg gca 334  
 Ala Leu Gly Gly Ser Ala Ile Tyr Gln Ser Thr Thr Asn Gln Ser Ala  
 20 25 30 35

aat aat agt cgt tca aat aca act agt aca aag gtt agt aac gtt tcg 382  
 Asn Asn Ser Arg Ser Asn Thr Thr Ser Thr Lys Val Ser Asn Val Ser  
 40 45 50

gta aat gtc aat acc gat gtt acc tct gca att gaa aaa gtt tca aat 430  
 Val Asn Val Asn Thr Asp Val Thr Ser Ala Ile Glu Lys Val Ser Asn  
 55 60 65

09869106-100901

tct gtc gtt tct gtt atg aat tat caa aaa gat aac tca caa agt agt Ser Val Val Ser Val Met Asn Tyr Gln Lys Asp Asn Ser Gln Ser Ser	478
70 75 80	
gac ttc agt tca att ttt ggt gga aat agc ggt tca agt tca tcg act Asp Phe Ser Ser Ile Phe Gly Gly Asn Ser Gly Ser Ser Ser Thr	526
85 90 95	
gat ggc tta cag ctt tct agt gaa ggc tct ggt gtc atc tac aaa aaa Asp Gly Leu Gln Leu Ser Ser Glu Gly Ser Gly Val Ile Tyr Lys Lys	574
100 105 110 115	
tct ggt ggt gat gcc tac gtt gta act aac tac cac gtt att gct ggt Ser Gly Gly Asp Ala Tyr Val Val Thr Asn Tyr His Val Ile Ala Gly	622
120 125 130	
aat agc tca ctt gat gtt ctg ctt tct ggt gga caa aaa gtc aaa gat Asn Ser Ser Leu Asp Val Leu Leu Ser Gly Gly Gln Lys Val Lys Asp	670
135 140 145	
tct gtg gtt ggt tat gat gaa tac aca gac ctt gct gtt ctt aaa atc Ser Val Val Gly Tyr Asp Glu Tyr Thr Asp Leu Ala Val Leu Lys Ile	718
150 155 160	
agt tct gaa cat gtc aaa gat gtg gcg aca ttc gct gat tct agt aaa Ser Ser Glu His Val Lys Asp Val Ala Thr Phe Ala Asp Ser Ser Lys	766
165 170 175	
tta aca att ggt gaa cct gcc att gcc gtt ggc tca cct tta ggt agt Leu Thr Ile Gly Glu Pro Ala Ile Ala Val Gly Ser Pro Leu Gly Ser	814
180 185 190 195	
caa ttt gca aac acc gca act gaa gga att tta tct gca aca agc cgt Gln Phe Ala Asn Thr Ala Thr Glu Gly Ile Leu Ser Ala Thr Ser Arg	862
200 205 210	
caa gtg act ttg acc caa gaa aat ggt caa aca act aat atc aat gca Gln Val Thr Leu Thr Gln Glu Asn Gly Gln Thr Thr Asn Ile Asn Ala	910
215 220 225	
att caa aca gat gct gcc att aac cct ggt aac tct gga ggg gct ttg Ile Gln Thr Asp Ala Ala Ile Asn Pro Gly Asn Ser Gly Gly Ala Leu	958
230 235 240	
att aat att gaa gga caa gtt att gga att act caa agt aaa att aca Ile Asn Ile Glu Gly Gln Val Ile Gly Ile Thr Gln Ser Lys Ile Thr	1006
245 250 255	
aca act gaa gat ggt tct act tct gtc gaa ggt tta gga ttt gcg att Thr Thr Glu Asp Gly Ser Thr Ser Val Glu Gly Leu Gly Phe Ala Ile	1054
260 265 270 275	
cct tct aat gat gtc gta aat atc att aat aaa ctt gaa gat gat ggt Pro Ser Asn Asp Val Val Asn Ile Ile Asn Lys Leu Glu Asp Asp Gly	1102
280 285 290	
aag att tca cgc cct gct tta ggt atc cga atg gtt gac ctt tca caa Lys Ile Ser Arg Pro Ala Leu Gly Ile Arg Met Val Asp Leu Ser Gln	1150
295 300 305	

```
<210> 2
<211> 408      -
<212> PRT
<213> Lactococcus lactis
```

```

<400> 2
Met Ala Lys Ala Asn Ile Gly Lys Leu Leu Leu Thr Gly Val Val Gly
  1          5          10          15
Gly Ala Ile Ala Leu Gly Gly Ser Ala Ile Tyr Gln Ser Thr Thr Asn
  20          25          30
Gln Ser Ala Asn Asn Ser Arg Ser Asn Thr Thr Ser Thr Lys Val Ser
  35          40          45
Asn Val Ser Val Asn Val Asn Thr Asp Val Thr Ser Ala Ile Glu Lys
  50          55          60

```



Val Ser Asn Ser Val Val Ser Val Met Asn Tyr Gln Lys Asp Asn Ser  
 65 70 75 80  
 Gln Ser Ser Asp Phe Ser Ser Ile Phe Gly Gly Asn Ser Gly Ser Ser  
 85 90 95  
 Ser Ser Thr Asp Gly Leu Gln Leu Ser Ser Glu Gly Ser Gly Val Ile  
 100 105 110  
 Tyr Lys Lys Ser Gly Gly Asp Ala Tyr Val Val Thr Asn Tyr His Val  
 115 120 125  
 Ile Ala Gly Asn Ser Ser Leu Asp Val Leu Leu Ser Gly Gly Gln Lys  
 130 135 140  
 Val Lys Asp Ser Val Val Gly Tyr Asp Glu Tyr Thr Asp Leu Ala Val  
 145 150 155 160  
 Leu Lys Ile Ser Ser Glu His Val Lys Asp Val Ala Thr Phe Ala Asp  
 165 170 175  
 Ser Ser Lys Leu Thr Ile Gly Glu Pro Ala Ile Ala Val Gly Ser Pro  
 180 185 190  
 Leu Gly Ser Gln Phe Ala Asn Thr Ala Thr Glu Gly Ile Leu Ser Ala  
 195 200 205  
 Thr Ser Arg Gln Val Thr Leu Thr Gln Glu Asn Gly Gln Thr Thr Asn  
 210 215 220  
 Ile Asn Ala Ile Gln Thr Asp Ala Ala Ile Asn Pro Gly Asn Ser Gly  
 225 230 235 240  
 Gly Ala Leu Ile Asn Ile Glu Gly Gln Val Ile Gly Ile Thr Gln Ser  
 245 250 255  
 Lys Ile Thr Thr Thr Glu Asp Gly Ser Thr Ser Val Glu Gly Leu Gly  
 260 265 270  
 Phe Ala Ile Pro Ser Asn Asp Val Val Asn Ile Ile Asn Lys Leu Glu  
 275 280 285  
 Asp Asp Gly Lys Ile Ser Arg Pro Ala Leu Gly Ile Arg Met Val Asp  
 290 295 300  
 Leu Ser Gln Leu Ser Thr Asn Asp Ser Ser Gln Leu Lys Leu Leu Ser  
 305 310 315 320  
 Ser Val Thr Gly Gly Val Val Val Tyr Ser Val Gln Ser Gly Leu Pro  
 325 330 335  
 Ala Ala Ser Ala Gly Leu Lys Ala Gly Asp Val Ile Thr Lys Val Gly  
 340 345 350  
 Asp Thr Ala Val Thr Ser Ser Thr Asp Leu Gln Ser Ala Leu Tyr Ser  
 355 360 365  
 His Asn Ile Asn Asp Thr Val Lys Val Thr Tyr Tyr Arg Asp Gly Lys  
 370 375 380

00659105-100901

Ser Asn Thr Ala Asp Val Lys Leu Ser Lys Ser Thr Ser Asp Leu Glu  
385 390 395 400

Thr Ser Ser Pro Ser Ser Ser Asn  
405

030305 100904  
T0600T 90163850